

## Antiplatelet Agents

Moscone West, 3rd Floor, Room 3016

Tuesday, October 29, 2013, 1:00 PM–3:15 PM

Abstract nos: 1-10

## TCT-1

## Do Proton Pump Inhibitors Interact With Clopidogrel? Insights From ADAPT-DES

Giora Weisz<sup>1</sup>, Ajay J. Kirtane<sup>2</sup>, Ke Xu<sup>3</sup>, Thomas Stuckey<sup>4</sup>, Bernhard Witzencbichler<sup>5</sup>, Michael Rinaldi<sup>6</sup>, Franz-Josef Neumann<sup>7</sup>, D. Christopher Metzger<sup>8</sup>, Timothy D. Henry<sup>9</sup>, David Cox<sup>10</sup>, Peter L. Duffy<sup>11</sup>, Bruce R. Brodie<sup>12</sup>, Ernest L. Mazzaferri<sup>13</sup>, Helen Parise<sup>14</sup>, Roxana Mehran<sup>15</sup>, Gregg Stone<sup>16</sup>

<sup>1</sup>Columbia University Medical Center and Cardiovascular Research Foundation, New York, United States, <sup>2</sup>Columbia University / Cardiovascular Research Foundation, New York, NY, <sup>3</sup>Cardiovascular Research Foundation, New York, NY, <sup>4</sup>Lebanon Cardiovascular Research Foundation, Greensboro, NC, <sup>5</sup>Charité Campus Benjamin Franklin, Berlin, Germany, <sup>6</sup>Sanger Heart & Vascular Institute, Charlotte, NC, <sup>7</sup>Universitäts-Herzzentrum Freiburg Bad Krozingen, Bad Krozingen, Germany, <sup>8</sup>Wellmont CVA Heart Institute, Kingsport, TN, <sup>9</sup>Minneapolis Heart Institute Foundation at Abbott Northwestern Hospital, Minneapolis, United States, <sup>10</sup>Lehigh Valley Health Network, Allentown, PA, <sup>11</sup>Pinehurst Cardiology, Pinehurst, NC, <sup>12</sup>LeBauer CV Research Foundation, Greensboro, NC, <sup>13</sup>Ohio State University, Dublin, OH, <sup>14</sup>Cardiovascular Research Foundation, New York, New York, <sup>15</sup>Mount Sinai Hospital, New York, NY, <sup>16</sup>Columbia University Medical Center and the Cardiovascular Research Foundation, New York, United States

**Background:** Selected proton pump inhibitors (PPI) interfere with clopidogrel metabolism, potentially attenuating P2Y<sub>12</sub> receptor-based platelet inhibition. Previous observational and randomized trials have reported varying results regarding the clinical significance of this pharmacologic interaction. We examined this relationship in the large-scale, prospective Assessment of Dual Anti-Platelet Therapy with Drug-Eluting Stents (ADAPT-DES) study.

**Methods:** Platelet reactivity testing was performed using the VerifyNow point-of-care assay in 8,583 patients at 11 US and German sites after successful DES implantation. All patients were treated with aspirin and clopidogrel, and were followed for 1 year. PPI were prescribed at the discretion of treating physicians.

**Results:** At the time of the post-procedure P2Y<sub>12</sub> test, 2,697 (31.4%) pts were on PPI, and 5,886 (68.6%) were not. Major baseline characteristics, P2Y<sub>12</sub> results and 1-year events are summarized in the Table. The use of PPI was an independent predictor of higher platelet reactivity (HPR) units (PRU) in a linear regression model ( $p < 0.0001$ ), and additionally was independently associated with HPR as defined as PRU  $> 208$  (OR 1.38 [1.25, 1.52],  $p = 0.0001$ ). At discharge, 2,163 (25.2%) pts were prescribed PPI, and 6,419 (74.8%) were not. In propensity-adjusted proportional hazards regression models, PPI use was independently associated with out of hospital mortality (HR 1.52 [1.09, 2.12],  $p = 0.01$ ) and MACE (HR 1.23 [1.00, 1.51],  $p = 0.049$ ).

Baseline Characteristics	PPI (n=2437)	No PPI (n=6146)	p-value
Mean age, years	64.4 ± 10.7	63.3 ± 10.8	<0.0001
Diabetes	34.8%	31.3%	0.002
Hypertension	83.7%	77.8%	<0.0001
Acute coronary syndrome	57.5%	49.0%	<0.0001
VerifyNow P2Y <sub>12</sub>			
Mean PRU	201.9 ± 97.3	181.6 ± 95.8	<0.0001
PRU > 208	49.3%	39.7%	<0.0001
%inhibition	36.0 ± 27.7	41.8 ± 28.3	<0.0001
1-Year Follow-up (out of hospital)			
Death	2.9%	1.5%	<0.0001
MI	2.1%	1.7%	0.26
MACE	6.9%	5.2%	0.004
Stent thrombosis (definite/probable)	1.1%	0.6%	0.05
Major Bleeding	5.3%	4.5%	0.11

**Conclusions:** In patients treated with clopidogrel after DES, the concomitant administration of PPI is associated with reduced platelet inhibition and adverse clinical outcomes. Additional studies are warranted to determine the risk-benefit ratio of PPI in patients after DES.

## TCT-2

## Comparative Effectiveness of Prasugrel vs. Clopidogrel Among Acute Myocardial Infarction Patients Treated With Percutaneous Coronary Intervention: 30-Day Outcomes from the TRANSLATE-ACS Observational Study

Tracy Y. Wang<sup>1</sup>, Marjorie Zettler<sup>2</sup>, Mark B. Effron<sup>3</sup>, Lisa McCoy<sup>4</sup>, Kevin Anstrom<sup>4</sup>, Gregg C. Fonarow<sup>5</sup>, John C. Messenger<sup>6</sup>, Gregg Stone<sup>7</sup>, David Cohen<sup>8</sup>, Timothy D. Henry<sup>9</sup>, Eric Peterson<sup>10</sup>


<sup>1</sup>Duke University, Durham, NC, <sup>2</sup>Lilly, Inc., Indianapolis, IN, <sup>3</sup>Eli Lilly and Company, Indianapolis, IN, <sup>4</sup>Duke Clinical Research Institute, Durham, NC, <sup>5</sup>UCLA Medical Center, Los Angeles, CA, <sup>6</sup>Denver VA Medical Center, Denver, CO, <sup>7</sup>Columbia University Medical Center and the Cardiovascular Research Foundation, New York, United States, <sup>8</sup>Saint Luke's Mid America Heart Institute, Kansas City, United States, <sup>9</sup>Minneapolis Heart Institute Foundation at Abbott Northwestern Hospital, Minneapolis, United States, <sup>10</sup>N/A, Durham, North Carolina

**Background:** Randomized trials have demonstrated the efficacy of prasugrel vs. clopidogrel in patients with acute MI, yet evidence of their effectiveness in real world practice is limited.

**Methods:** We studied use of prasugrel vs. clopidogrel among 11,417 STEMI and NSTEMI patients treated with PCI in the TRANSLATE-ACS study from 4/2010 to 10/2012. We used multivariable Cox models to compare 30-day MACE (death, recurrent MI, stroke, or unplanned revascularization) and any GUSTO defined bleeding.

**Results:** Prasugrel was used in 2,997 MI patients (26%) during PCI. Patients treated with prasugrel were younger (median 57 vs. 61 yrs), more likely to present with STEMI (59% vs. 49%), and less likely to have prior MI (15% vs. 21%) or diabetes (24% vs. 27%) than those receiving clopidogrel ( $p < 0.01$  for all). Prasugrel was used in 57 (9%) of patients with prior stroke/TIA, 64 (5%) over age 75, and 74 (14%) of patients  $< 60$  kg. Compared with clopidogrel, prasugrel was more often started during/after PCI, and used with bivalirudin or GIIb/IIIa inhibitor. Unadjusted curves for 30-day MACE and bleeding diverged, but were not significantly different between prasugrel and clopidogrel treated patients (Figure). Multivariable analyses did not demonstrate significant differences in MACE (HR 0.96, 95%CI 0.79, 1.16) and bleeding (HR 1.04, 95%CI 0.77, 1.39).

**Conclusions:** While differences exist in patients receiving these drugs, the 30-day effectiveness and safety of prasugrel vs. clopidogrel were not significantly different in routine practice. Long-term outcomes comparisons are necessary and ongoing.



## TCT-3

## Outcomes in STEMI patients treated with clopidogrel or prasugrel: A propensity adjusted analysis from the INFUSE-AMI Trial

Keith G. Oldroyd<sup>1</sup>, Sorin Brenner<sup>2</sup>, Akiko Maehara<sup>3</sup>, Magdi El-Omar<sup>4</sup>, Bernhard Witzencbichler<sup>5</sup>, Ke Xu<sup>6</sup>, Roxana Mehran<sup>7</sup>, C. Michael Gibson<sup>8</sup>, Gregg W. Stone<sup>9</sup>

<sup>1</sup>University of Glasgow, Glasgow, United Kingdom, <sup>2</sup>New York Methodist Hospital, Brooklyn, New York, NY, <sup>3</sup>Cardiovascular Research Foundation and Columbia University Medical Center, New York, United States, <sup>4</sup>Manchester Heart Centre, Manchester, United Kingdom, <sup>5</sup>Charité Campus Benjamin Franklin, Berlin, Germany, <sup>6</sup>Cardiovascular Research Foundation, New York, NY, <sup>7</sup>Mount Sinai Hospital, New York, NY, <sup>8</sup>Beth Israel Deaconess Med Ctr - Harvard Medical School, Boston, MA, <sup>9</sup>Cardiovascular Research Foundation, NY, NY

**Background:** Prasugrel was superior to Clopidogrel in a large trial of acute coronary syndromes patients undergoing percutaneous coronary intervention (PCI) with heparin-based anticoagulation. It is not known whether more potent platelet inhibition with Prasugrel rather than Clopidogrel affects infarct size and clinical outcomes when primary PCI is performed with bivalirudin anticoagulation.